

EDITORIAL COMMENT

## Aspiration Thrombectomy

### It's About Time\*

Prakash Balan, MD, H. Vernon Anderson, MD

Houston, Texas

In an elegant meta-analysis published 5 years ago, Bavry et al. (1) reported that aspiration thrombectomy prior to percutaneous coronary intervention (PCI) in acute ST segment-elevation myocardial infarction (STEMI) can be beneficial. Using data from 30 studies of 6,415 patients, they found improvements in 2 surrogate markers of coronary flow and reperfusion, with more Thrombolysis In Myocardial Infarction (TIMI) blush grade of 3 at the conclusion of the procedure, and more ST-segment resolution (STR) in the group with aspiration thrombectomy compared with PCI without thrombectomy. More important, at a mean follow-up interval of 5 months, mortality was 2.7% in the aspiration thrombectomy group compared with 4.4% in the PCI-only

See page 1409

group ( $p = 0.018$ ). Neither mechanical thrombectomy nor embolic protection devices were found to provide similar benefits or any benefit at all in the meta-analysis. Subsequently, the American College of Cardiology/American Heart Association (ACC/AHA) STEMI guidelines endorsed the use of aspiration thrombectomy with PCI in STEMI, with a Class IIa (Level of Evidence: B) recommendation (2). The recently updated 2013 ACCF/AHA STEMI guidelines retain this endorsement (3). Nonetheless, the use of aspiration thrombectomy during PCI in STEMI has remained controversial. On one hand, removal of some or possibly most of the offending thrombus at the culprit lesion site may prevent distal embolization, improve epicardial coronary flow, and reduce microvascular obstruction and the no-reflow phenomenon. On the other hand, the thrombectomy catheter itself is one more device that may cause distal thrombus embolization, and its insertion adds procedure time prior to definitive balloon angioplasty and/or stenting. Not everyone is convinced there is benefit. Partly this is because there has been a great deal of heterogeneity among individual clinical trials, most of which have been


small, as was documented in the meta-analysis (1). The recently reported TROFI (Thrombus Aspiration on Flow Area in Patients With ST-Elevation Myocardial Infarction) trial (4) is likely not of much help either. Using optical frequency domain imaging, the TROFI investigators found in 141 STEMI patients that minimal flow area at the lesion site was no different when aspiration thrombectomy was performed prior to stenting compared with stenting alone. What that finding might mean is unknown at present. Myocardial blush grades, STR, and clinical events also were not different, but the TROFI trial was not designed or powered for those secondary outcomes.

Meanwhile, in this issue of the *Journal*, Kumbhani et al. (5) of the previous meta-analysis have now updated their work with several important additions. A completely new meta-analysis was done, and this one included 5,534 patients from 25 trials. In addition to TIMI blush grades, STR, and intermediate-term mortality, the authors also examined data on infarct size, left ventricular function (ejection fraction; LVEF), and total ischemic time, hoping to gain insight into mechanisms of benefit. Similar to their previous findings, both TIMI blush grade 3 and STR were significantly better with aspiration thrombectomy compared with PCI only, and mortality at 6 months was lower (2.7% vs. 3.9%;  $p = 0.049$ ). Mechanical thrombectomy again did not exhibit benefit compared with PCI alone. The data extracted regarding infarct size and LVEF are interesting but not conclusive, and this further reinforces the need for larger randomized trials that address these clinical measures directly using uniform methods. In the trials included in this meta-analysis, the follow-up assessments of infarct size and LVEF were done at a mean of 36 days (range: from hospital discharge to 6 months) and involved a variety of methods, including echocardiography, radionuclide perfusion studies, and magnetic resonance imaging. Selection bias, survival bias, and incomplete follow-up no doubt interfered with the comparisons, and it is not surprising that there were no differences in infarct size (mean: 17.1% vs. 17.3%;  $p = 0.64$ ) or LVEF (mean: 53.0% vs. 52.8%;  $p = 0.32$ ). Also, these relatively smaller infarcts and essentially normal LVEFs, coupled with the relatively lower mortalities, all suggest that stringent clinical selection allowed only less clinically ill patients into the trials. Again, this is no surprise.

Finally, it is important and noteworthy that these authors extracted ischemic times, which they are careful to define as *total ischemic time*, generally taken to be the time from onset of infarct symptoms until device activation. Mean ischemic times in these studies ranged from 2.4 h to 7.6 h. We and others have shown that total ischemic time is the correct focus of attention for reperfusion in STEMI (6,7). To further illustrate its importance, we used the ischemic time and infarct size data reported in the present meta-analysis (Fig. 1). Based on data from studies in animals and humans, it appears that a total ischemic time of <120 min is a worthwhile and achievable goal to keep infarct size as small as possible and mortality as low as possible. The present meta-analysis data do not contradict this. In conclusion, the proper

\*Editorials published in the *Journal of the American College of Cardiology* reflect the views of the authors and do not necessarily represent the views of JACC or the American College of Cardiology.

From the Division of Cardiology, University of Texas Health Science Center, Houston, Texas. Both authors have reported that they have no relationships relevant to the contents of this paper to disclose.



**Figure 1** Ischemic Time Versus Infarct Size

The **x-axis** shows mean ischemic time in hours. The **y-axis** shows infarct size as a percentage of left ventricle. A simple regression line is shown; the data points are nonadjusted and nonweighted. Data from Tables 1 and 2 in the paper by Kumbhani et al. (5).

interpretation of this newly updated meta-analysis is that, like its predecessor, it provides solid, if indirect, evidence that aspiration thrombectomy prior to PCI in STEMI is useful and, like all infarct treatments, should be applied as early as possible, ideally within 120 min of ischemic time. We agree with the authors that until larger clinical trials are done, the present ACCF/AHA STEMI guideline recommendation (Class IIa, Level of Evidence: B) remains appropriate.

**Reprint requests and correspondence:** Dr. H. Vernon Anderson, Cardiology Division–UTHSCH, 6431 Fannin, Room 1.246, Houston, Texas 77030. E-mail: [h.v.anderson@uth.tmc.edu](mailto:h.v.anderson@uth.tmc.edu).

## REFERENCES

1. Bavry AA, Kumbhani DJ, Bhatt DL. Role of adjunctive thrombectomy and embolic protection devices in acute myocardial infarction: a comprehensive meta-analysis of randomized trials. *Eur Heart J* 2008; 29:2989–3001.
2. Kushner FG, Hand M, Smith SC Jr., et al. 2009 focused updates: ACC/AHA guidelines for the management of patients with ST-elevation myocardial infarction (updating the 2004 guideline and 2007 focused update) and ACC/AHA/SCAI guidelines on percutaneous coronary intervention (updating the 2005 guideline and 2007 focused update): a report of the American College of Cardiology Foundation/American Heart Association Task Force on Practice Guidelines. *J Am Coll Cardiol* 2009;54:2205–41.
3. O’Gara PT, Kushner FG, Ascheim DD, et al. 2013 ACCF/AHA guideline for the management of ST-elevation myocardial infarction: executive summary: a report of the American College of Cardiology Foundation/American Heart Association Task Force on Practice Guidelines. *J Am Coll Cardiol* 2013;61: 485–510.
4. Yoshinobu Y, Thuesen L, van Geuns RJ, et al. Randomized study to assess the effect of thrombus aspiration on flow area in patients with ST-elevation myocardial infarction: an optical frequency domain imaging study—TROFI trial. *Eur Heart J* 2013;34: 1050–60.
5. Kumbhani DJ, Bavry AA, Desai MY, Bangalore S, Bhatt DL. Role of aspiration and mechanical thrombectomy in patients with acute myocardial infarction undergoing primary angioplasty: an updated meta-analysis of randomized trials. *J Am Coll Cardiol* 2013;62: 1409–18.
6. Denktas AE, Anderson HV, McCarthy J, Smalling RW. Total ischemic time: the correct focus of attention for optimal ST-segment elevation myocardial infarction care. *J Am Coll Cardiol Intv* 2011;4: 599–604.
7. Reyes MM, Gunter AL, Daraban N, et al. Ischemic time: a better predictor than door-to-balloon time for infarct size in STEMI by cardiac magnetic resonance imaging (abstr). *Circulation* 2011;124: A15506.

**Key Words:** meta-analysis ■ mortality myocardial infarction ■ outcomes ■ thrombectomy.